

# The character and the wave front set correspondence in the stable range

Tomasz Przebinda

University of Oklahoma  
Norman, OK, USA

**50th Seminar Sophus Lie**  
Bedlewo, Poland, September 25 - October 1, 2016

# Howe's Theory of Rank

$$E = \begin{pmatrix} 0 & & 1 \\ & \ddots & \\ 1 & & 0 \end{pmatrix}.$$

$$\mathrm{Sp}_{2n} = \left\{ g \in \mathrm{M}_{2n}(\mathbb{R}); g^t \begin{pmatrix} 0 & E \\ -E & 0 \end{pmatrix} g = \begin{pmatrix} 0 & E \\ -E & 0 \end{pmatrix} \right\}$$

$$\mathrm{P} = \left\{ \begin{pmatrix} a & c \\ 0 & {}^t a^{-1} \end{pmatrix}; a \in \mathrm{GL}_n, c = {}^t c \in \mathrm{M}_n \right\}$$

$$\mathrm{N} = \left\{ \begin{pmatrix} I & c \\ 0 & I \end{pmatrix}; c = {}^t c \in \mathrm{M}_n \right\}$$

$$\hat{\mathrm{N}} = \{ \hat{c} = {}^t \hat{c} \in \mathrm{M}_n \}, \quad \hat{c} \left( \begin{pmatrix} I & c \\ 0 & I \end{pmatrix} \right) = \exp(2\pi i \mathrm{tr}(\hat{c}c)).$$

Suppose

$\Pi$  is an irreducible unitary representation of a cover of  $Sp_{2n}$   
and  
 $\Pi|_{\mathbb{N}}$  is supported on matrices of maximal rank  $r < n$ .

Then

$\Pi$  factors through a double cover

$$\widetilde{Sp}_{2n} \rightarrow Sp_{2n}$$

of the symplectic group, which is trivial if  $r$  is even and is equal to the metaplectic cover if  $r$  is odd. Moreover...

$$\begin{aligned}
 P_1 &= \left\{ \left( \begin{array}{ccc} a_1 & b_1 & c_1 \\ 0 & a_2 & b_1^* \\ 0 & 0 & {}^t a_1^{-1} \end{array} \right) \in \mathrm{Sp}_{2n}; a_1 \in \mathrm{GL}_r, a_2 \in \mathrm{Sp}_{2n-2r} \right\} \\
 &= (\mathrm{GL}_r \times \mathrm{Sp}_{2n-2r}) \cdot N_1
 \end{aligned}$$

There are unique integers  $p \geq q \geq 0$  with  $p + q = r$ , an irreducible unitary representation  $\Pi'_O$  of  $O_{p,q}$  and a character  $\alpha : \tilde{O}_{p,q} \rightarrow \mathbb{C}$  such that

$$\Pi|_{\tilde{P}_1} = \mathrm{Ind}_{(O_{p,q} \times \mathrm{Sp}_{2n-2r}) \cdot N_1}^{\tilde{P}_1} ((\alpha \Pi'_O \otimes 1) \otimes \omega_{p,q}).$$

There is a subset  $(\widehat{\mathrm{Sp}}_{2n})_{p,q} \subseteq \widehat{\mathrm{Sp}}_{2n}$  such that the above formula gives a bijection

$$(\widehat{\mathrm{Sp}}_{2n})_{p,q} \ni \Pi \longleftrightarrow \Pi'_O \in \hat{O}_{p,q}.$$

E.g. the rank one representations ( $r = 1$ ) are the two Weil representations of the metaplectic group.

# The Metaplectic Group

$(W, \langle \cdot, \cdot \rangle)$ ;  $\mathrm{Sp}, \mathfrak{sp} \subseteq \mathrm{End}(W)$ ;  $J \in \mathfrak{sp}$ ,  $J^2 = -I$ ,  $\langle J\cdot, \cdot \rangle > 0$ ;  $g, g_1, g_2 \in \mathrm{Sp}$ ;  
 $\chi(r) = e^{2\pi i r}$ ,  $r \in \mathbb{R}$ .

The restriction of  $J_g = J^{-1}(g - 1)$  to  $J_g W$  is invertible. Let

$$C(g_1, g_2) = \sqrt{\left| \frac{\det(J_{g_1})_{J_{g_1}W} \det(J_{g_2})_{J_{g_2}W}}{\det(J_{g_1 g_2})_{J_{g_1 g_2}W}} \right|} \chi\left(\frac{1}{8} \mathrm{sgn}(q_{g_1, g_2})\right),$$

where  $\mathrm{sgn}(q_{g_1, g_2})$  is the signature of the symmetric form

$$\begin{aligned} q_{g_1, g_2}(u', u'') &= \frac{1}{2} \langle (g_1 + 1)(g_1 - 1)^{-1} u', u'' \rangle \\ &+ \frac{1}{2} \langle (g_2 + 1)(g_2 - 1)^{-1} u', u'' \rangle \\ &(u', u'' \in (g_1 - 1)W \cap (g_2 - 1)W). \end{aligned}$$

$$\begin{aligned} \widetilde{\mathrm{Sp}} &= \left\{ \tilde{g} = (g, \xi) \in \mathrm{Sp} \times \mathbb{C}, \quad \xi^2 = i^{\dim(g-1)W} \det(J_g)_{J_g W}^{-1} \right\}, \\ &(g_1, \xi_1)(g_2, \xi_2) = (g_1 g_2, \xi_1 \xi_2 C(g_1, g_2)). \end{aligned}$$

# The Weil Representation

$$W = X \oplus Y; \text{Op} : \mathcal{S}^*(X \times X) \rightarrow \text{Hom}(\mathcal{S}(X), \mathcal{S}^*(X))$$

$$\text{Op}(K)v(x) = \int_X K(x, x')v(x') dx'.$$

$$\text{Weyl transform } \mathcal{K} : \mathcal{S}^*(W) \rightarrow \mathcal{S}^*(X \times X)$$

$$\mathcal{K}(f)(x, x') = \int_Y f(x - x' + y)\chi\left(\frac{1}{2}\langle y, x + x' \rangle\right) dy.$$

A singular imaginary Gaussian

$$\chi_{c(g)}(u) = \chi\left(\frac{1}{4}\langle (g+1)(g-1)^{-1}u, u \rangle\right) \quad (u = (g-1)w, w \in W).$$

For  $\tilde{g} = (g, \xi) \in \widetilde{\text{Sp}}$  define

$$\Theta(\tilde{g}) = \xi, \quad T(\tilde{g}) = \Theta(\tilde{g})\chi_{c(g)}\mu_{(g-1)W}, \quad \omega(\tilde{g}) = \text{Op} \circ \mathcal{K} \circ T(\tilde{g}),$$

where  $\mu_{(g-1)W}$  is the Lebesgue measure on the subspace  $(g-1)W$  normalized so that the volume of the unit cube with respect to the form  $\langle J \cdot, \cdot \rangle$  is 1. In these terms,  $(\omega, L^2(X))$  is the Weil representation of  $\widetilde{\text{Sp}}$  attached to the character  $\chi$ .

## Dual Pairs

Subgroups  $G, G' \subseteq \mathrm{Sp}(W)$  act reductively on  $W$ ,  $G'$  is the centralizer of  $G$  in  $\mathrm{Sp}$  and  $G$  is the centralizer of  $G'$  in  $\mathrm{Sp}$ .

$G, G'$	stable range
$\mathrm{GL}_n(\mathbb{D}), \mathrm{GL}_m(\mathbb{D})$	$n \geq 2m$
$\mathrm{O}_{p,q}, \mathrm{Sp}_{2n}(\mathbb{R})$	$p, q \geq 2n$
$\mathrm{Sp}_{2n}(\mathbb{R}), \mathrm{O}_{p,q}$	$n \geq p + q$
$\mathrm{O}_p(\mathbb{C}), \mathrm{Sp}_{2n}(\mathbb{C})$	$p \geq 4n$
$\mathrm{Sp}_{2n}(\mathbb{C}), \mathrm{O}_p(\mathbb{C})$	$n \geq p$
$\mathrm{U}_{p,q}, \mathrm{U}_{r,s}$	$p, q \geq r + s$
$\mathrm{Sp}_{p,q}, \mathrm{O}_{2n}^*$	$p, q \geq n$
$\mathrm{O}_{2n}^*, \mathrm{Sp}_{p,q}$	$n \geq 2(p + q)$

## Howe's Correspondence

$\mathcal{R}(\tilde{G}, \omega) \subseteq \mathcal{R}(\tilde{G})$  irreducible admissible representations realized as quotients of  $\mathcal{S}(X)$  by closed  $\tilde{G}$ -invariant subspaces.  $\Pi \in \mathcal{R}(\tilde{G}, \omega)$ ,  $N_\Pi \subseteq \mathcal{S}(X)$  the intersection of all the closed  $G$ -invariant subspaces  $N \subseteq \mathcal{S}(X)$  such that  $\Pi$  is infinitesimally equivalent to  $\mathcal{S}(X)/N$ . This is a representation of both  $\tilde{G}$  and  $\tilde{G}'$ . It is infinitesimally isomorphic to

$$\Pi \otimes \Pi'_1,$$

for some representation  $\Pi'_1$  of  $\tilde{G}'$ . Howe proved that  $\Pi'_1$  is a finitely generated admissible quasisimple representation of  $\tilde{G}'$ , which has a unique irreducible quotient  $\Pi' \in \mathcal{R}(\tilde{G}', \omega)$ . Conversely, starting with  $\Pi' \in \mathcal{R}(\tilde{G}', \omega)$  and applying the above procedure with the roles of  $G$  and  $G'$  reversed, we arrive at the representation  $\Pi \in \mathcal{R}(\tilde{G}, \omega)$ . The resulting bijection

$$\mathcal{R}(\tilde{G}, \omega) \ni \Pi \longleftrightarrow \Pi' \in \mathcal{R}(\tilde{G}', \omega)$$

is called Howe's correspondence, or local  $\theta$  correspondence, for the pair  $G, G'$ .



Let  $(G, G')$  be a dual pair in the stable range with  $G'$  - the smaller member. If  $\Pi'$  is unitary then  $\Pi$  is unitary. Howe's theory of rank is compatible with Howe's correspondence. In particular for the pair  $(\mathrm{Sp}_{2n}(\mathbb{R}), \mathrm{O}_{p,q})$  with  $p + q < n$ , there is a character  $\alpha'$  of the preimage of the orthogonal group in the metaplectic group such that

$$\Pi' = \alpha' \Pi'_O.$$

In particular  $\Pi|_{\tilde{\mathbb{P}}_1}$  is understood in terms of  $\Pi'$ .

An explicit description of all the  $\Pi(g)$ ,  $g \in \tilde{G}$ , seems out of reach. Instead one may try to **describe the distribution character  $\Theta_\Pi$  in terms of  $\Theta_{\Pi'}$** .

# The wave front set of a distribution

The wave front set of a distribution  $u$  on an Euclidean space  $V$  at a point  $v \in V$ , denoted  $WF_v(u)$  is the complement of the set of all pairs  $(v, v^*)$ ,  $v^* \in V^*$ , such that there is a test function  $\phi \in C_c^\infty(V)$  with  $\phi(v) \neq 0$  and an open cone  $\Gamma \subseteq V^*$  containing  $v^*$  such that

$$|\mathcal{F}(\phi u)(v_1^*)| \leq C_N(1 + |v_1^*|)^{-N} \quad (v_1^* \in \Gamma, N = 0, 1, 2, \dots).$$

This notion behaves well under diffeomorphisms, so one may define  $WF(u) \subseteq T^*M$  as the union of the wave front sets at the individual points for any distribution  $u$  on a manifold  $M$ .

# The Cauchy Harish-Chandra Integral

Assume that the rank of  $G'$  is smaller or equal than the rank of  $G$ .  
 $\theta$ -invariant Cartan subgroup  $H' \subseteq G'$ ;  $A' = \{h \in H'; \theta(h) = h^{-1}\}$ ;  
 $A'' \subseteq \text{Sp}$  centralizer of  $A'$ ,  $A''' \subseteq \text{Sp}$  centralizer of  $A''$ ;  $d\dot{w}$  measure on  
 $A''' \backslash W$  defined by

$$\int_W \phi(w) dw = \int_{A''' \backslash W} \int_{A'''} \phi(aw) da d\dot{w}.$$

$$\text{Chc}(f) = \int_{A''' \backslash W} \int_{\tilde{A}''} f(g) T(g)(w) dg d\dot{w} \quad (f \in C_c^\infty(\tilde{A}'')).$$

For any  $h \in H'^{\text{reg}}$ , the intersection of the wave front set of the distribution  $\text{Chc}$  with the conormal bundle of the embedding

$$\tilde{G} \ni \tilde{g} \longrightarrow \tilde{h}\tilde{g} \in \tilde{A}''$$

is empty. Hence there is a unique restriction of the distribution  $\text{Chc}$  to  $\tilde{G}$ , denoted  $\text{Chc}_{\tilde{h}}$ .

## The distribution $\Theta'_{\Pi'}$

Set  $D(h) = |\det(\text{Ad}(h^{-1}) - 1)_{\mathfrak{g}'/\mathfrak{h}'}|$ .

Weyl integration formula

$$\int_{\tilde{G}'} \phi(g) dg = \sum \frac{1}{|W(H')|} \int_{\tilde{H}'^{reg}} D(h) \int_{\tilde{G}'/\tilde{H}'} \phi(g\tilde{h}g^{-1}) dg d\tilde{h},$$

For  $f \in C_c^\infty(\tilde{G})$  define

$$\Theta'_{\Pi'}(f) = C_{\Pi'} \sum \frac{1}{|W(H')|} \int_{\tilde{H}'^{reg}} D(h) \Theta_{\Pi'}(\tilde{h}^{-1}) \text{Ch}_{c_{\tilde{h}}}(f) d\tilde{h}.$$

This is an invariant eigen-distribution on  $\tilde{G}$  with the correct infinitesimal character.

# The equality $\Theta'_{\Pi'} = \Theta_{\Pi}$

## Theorem

*Let  $(G, G')$  be a dual pair in the stable range with  $G'$  - the smaller member. If  $\Pi'$  is unitary then  $\Theta'_{\Pi'} = \Theta_{\Pi}$ .*

**Idea of the proof.** We show that the two distributions are equal on a Zariski open subset  $\tilde{G}'' \subseteq \tilde{G}$ . Since both  $\Theta_{\Pi}$  and  $\Theta'_{\Pi'}$  is an invariant eigendistribution, Harish-Chandra Regularity Theorem implies that they are equal everywhere.

# The moment maps

The maps  $\tau_{\mathfrak{g}} : W \rightarrow \mathfrak{g}^*$  and  $\tau_{\mathfrak{g}'^*} : W \rightarrow \mathfrak{g}'^*$  defined by

$$\tau_{\mathfrak{g}}(z) = \langle z(w), w \rangle \quad (z \in \mathfrak{g}, w \in W)$$

(and similarly for  $\mathfrak{g}'$ ) occur naturally in the structure of the Weil representation.

If  $\mathcal{N}(\mathfrak{g}^*) \subseteq \mathfrak{g}^*$  denotes the subset of the nilpotent elements, then

$$\tau_{\mathfrak{g}}(\tau_{\mathfrak{g}'^*}^{-1}(\mathcal{N}(\mathfrak{g}'^*))) \subseteq \mathcal{N}(\mathfrak{g}^*).$$

The equality  $WF(\Pi) = \tau_{\mathfrak{g}}(\tau_{\mathfrak{g}'}^{-1}(WF(\Pi')))$

$$WF(\Pi) = WF_1(\Theta_{\Pi}) \subseteq \mathcal{N}(\mathfrak{g}^*).$$







## Theorem

*Let  $(G, G')$  be a dual pair in the stable range with  $G'$  - the smaller member. If  $\Pi'$  is unitary then  $WF(\Pi) = \tau_{\mathfrak{g}}(\tau_{\mathfrak{g}'}^{-1}(WF(\Pi')))$ .*







This follows from the equality  $\Theta'_{\Pi'} = \Theta_{\Pi}$  and the properties of the distribution  $\Theta'_{\Pi'}$ , except when  $G' \neq G'_1$  (the Zariski identity component of  $G'$ ), the restriction of  $\Pi'$  to  $\tilde{G}'_1$  is irreducible and  $WF(\Pi')$  contains more than one orbit of maximal dimension. In that last case, we use the result of Loke and Ma, who computed the correspondence of the associated varieties of the Harish-Chandra modules of  $\Pi$  and  $\Pi'$  and a theorem of Schmid and Vilonen proving a conjecture of Barbasch and Vogan.

**Remark:** another possibility would be to extend the notion of the Cauchy Harish-Chandra integral to the full group  $\tilde{G}'$ , not just  $\tilde{G}'_1$ .

# References

-  R. HOWE: *Small unitary representations of classical groups*. Group representations, ergodic theory, operator algebras, and mathematical physics (Berkeley, Calif., 1984), 121-150
-  A.-M. AUBERT AND T. PRZEBINDA: *A reverse engineering approach to the Weil Representation*. CEJM, **12** (2014), 1200–1285
-  R. HOWE: *Transcending Classical Invariant Theory*. J. Amer. Math. Soc., **2** (1989), 535-552
-  JIAN-SHU LI: *Singular unitary representations of classical groups*. Inven. Math., **97** (1989), 237-255
-  JIAN-SHU LI: *On the classification of irreducible low rank unitary representations of classical groups*. Compos. Math., **71** (1989), 29-48
-  L. HÖRMANDER *The Analysis of Linear Partial Differential Operators I*. Springer Verlag, (1983)



-  T. PRZEBINDA: *A Cauchy Harish-Chandra Integral, for a real reductive dual pair*. Inven. Math., **141** (2000), 299–363
-  R. HOWE: *Wave Front Sets of Representations of Lie Groups*. Automorphic forms, Representation Theory and Arithmetic, Tata Institute of Fundamental Research, Bombay (1981), 117-140
-  F. BERNON AND T. PRZEBINDA: *The Cauchy Harish-Chandra integral and the invariant eigendistributions*. International Mathematics Research Notices, **14** (2014) 3818–3862
-  T. PRZEBINDA: *The character and the wave front set correspondence in the stable range*. Arxiv: 1602.08401, 19 pages
-  H. Y. LOKE AND J.J. MA: *Invariants and  $K$ -spectrum of local theta lifts*. Compos. Math., **151** (2015), 179-206
-  W. SCHMID AND K. VILONEN: *Characteristic cycles and wave front cycles of representations of reductive Lie groups*. Ann. of Math. (2), **151** (2000), 1071 – 1118

# Thank you